

Giant pulses in Millisecond Pulsars

B. C. Joshi

National Center for Radio Astrophysics, Ganeshkhind, Pune, India

M. Kramer, A. G. Lyne, M. McLaughlin

Jodrell Bank Observatory, University of Manchester, Macclesfield, UK

I. H. Stairs

*Dept. of Physics and Astronomy, University of British Columbia, 6224
 Agricultural Road, Vancouver, B.C. V6T 1Z1 Canada*

Abstract. Giant pulses (GPs), occasional individual pulses with an intensity 100 times the average intensity, have been detected in four pulsars todate. Their origin is not well understood, but studies suggest a connection between the strength of magnetic field at the light cylinder B_{lc} and the existence of GPs. Here, we report on detection of significant Large Amplitude Pulses (LAPs) in two more pulsars with high values of B_{lc} , PSRs J0218+4232 and B1957+20, observed using Giant Meterwave Radio Telescope (GMRT).

1. Introduction

Giant pulses (GPs) have been reported in four pulsars (B0531+21, B1937+21, B1821–24 and B0540–69) todate (Staelin & Reifenstein 1968; Lundgren et al. 1995; Cognard et al. 1996; Romani & Johnston 2001; Johnston & Romani 2003). Three of these pulsars are millisecond pulsars (MSPs) and also show strongly pulsed hard X-ray profiles (Takahashi et al. 2001). The radio GPs occur in a narrow phase window close to the high energy non-thermal pulse indicating a common magnetospheric origin. All these pulsars have a high value of B_{lc} . We have used GMRT to search for GPs in candidate MSPs with a non-thermal high energy emission and a range of B_{lc} and report detection of such pulses in two more pulsars, PSR J0218+4232 and B1957+20.

We obtained about 3600 s of data on each pulsar using 20 to 22 GMRT antennae in an incoherent mode at 610 MHz with 16 MHz bandwidth. The expected RMS noise in the above configuration of GMRT for a sampling time of 258 μ s is about 1 Jy. The data in two subbands (8 MHz each) were dedispersed to a common sky frequency (610 MHz). The periods with a peak greater than 3.5 times RMS in both bands at the same sample were identified as Large amplitude pulses (LAPs). This procedure disperses any narrow interference spike discriminating against interference. A number of marginal LAPs, i.e. pulses with a peak between 3.0 to 3.5 times RMS, were also identified.




Figure 1. LAPs in PSR J0218+4232 and B1957+20. a) The left panel shows the baseline subtracted radio integrated profile for PSR J0218+4232 with the LAPs marked with the filled circles at the phase of their occurrence. The energy in the LAP is labeled on the right. b) The right panel shows the GP detected in PSR B1957+20 after removing the off-pulse mean.

2. Results

We searched 2.2 million periods for PSR J0218+4232 ($B_{lc} \sim 3.2 \times 10^5$ G) and found three significant LAPs. Figure 1a shows the integrated profile for this pulsar and the detected LAPs are marked with filled circles. The largest of these had an intensity 51 times the mean intensity (Intensity of LAP, $I_{lap} \sim 1341$ Jy- μ s). Our data also consisted of 9 marginal LAPs, 7 of which occur between phase 0.89 - 1.2, which is the phase interval corresponding to one of the high energy peaks. We searched about 1 million periods for PSR B1957+20 ($B_{lc} \sim 3.8 \times 10^5$ G) and found one significant LAP, shown in Figure 1b, with an intensity 129 times the mean intensity ($I_{lap} \sim 925$ Jy- μ s). In addition, we also detected 5 marginal LAPs.

PSRs B1957+20 and J0218+4232 have the fourth and sixth highest values of B_{lc} respectively of known radio pulsars. Hence, these new detections support a connection between the magnitude of B_{lc} and the existence of GPs and LAPs. The data for pulsars with B_{lc} marginally below 10^5 G are being analyzed currently.

References

- Cognard, I. et al. 1996, ApJ, 457, 81
- Johnston, S., & Romani, R. W. 2003, private communication
- Lundgren, S. C. et al. 1995, ApJ, 453, 433
- Romani, R. W., & Johnston, S. 2001, ApJ, 557, L93
- Staelin, D. H., & Reifenstein, E. C. 1968, Science, 162, 1481
- Takahashi, M. et al. 2001, ApJ, 554, 316